



- **OXYPY®** is a bio-insecticide for internal use effective on fleas and bedbugs.
- **OXYPY®** is a ready-for-use solution, conceived to be used exclusively with the diffusion appliances of the **NOCOTECH®** range or in local applications with a spray.

REFERENCES AND PACKAGING

Reference	Packaging
4230.001	1 Liter
4230.020	20 Litre Canister

COMPOSITION

S-methopren 0.5% • EC=613-834-0 / CAS=65733-16-6.

Pyrethrin 2% • EC=232-319-8 / CAS=8003-34-7.

Pyperonyl butoxide 5% • EC=200-076-7 / CAS=51-03-6.

STORAGE

- Store the product in the original packaging, vertically and in a cool and well ventilated place.
- Shelf-life : In the closed original packaging : 2 years from manufacturing date.
Once opened : 2 months from opening date.

OXYPHARM

829 rue Marcel Paul
94500 Champigny-sur-Marne
commercial@oxypharm.net
T: +33 1 45 18 78 70

www.oxypharm.net



PRECAUTION FOR USE

- Refer to the material safety data sheet, available on request by email: commercial@oxypharm.net.

INSTRUCTIONS FOR USE

Protocol of use

- Follow the instructions for use of the diffusion appliance of the range NOCOTECH® (cf. user's manual and quickstart document).
- Shake the bottle before each use.
- Attach the 1L bottle to the diffusion appliance NOCOSPRAY® or the 20L tank on the diffusion appliance NOCOMAX®.
- Remove the mattress off the bed and display all parts with difficult access.
- Clean with a steam system (if possible) all fabrics, woods and structures where bedbugs eggs might settle and vacuum with a vacuum cleaner.
- Apply the product in all the nooks of the bed, using the provided spray head.
- Leave the mattress on the edge and begin the treatment of the room.
- On the device, set the volume (V) of the room to be treated multiplied by 2.
As an example: a 20m² room with a height of approximately 2,50m will have a volume of 20 x 2,50m = 50m³.
The device will have to be set on 100m³.
- After the end of diffusion, respect a dwell time of 2 hours minimum and then ventilate the room (opening the windows) during 10 minutes.
- After the treatment, remove the bottle from the appliance and close it.
- It is advised to carry out overhaul treatments every 6 weeks.

Overhaul treatment

Follow the same instruction than for the first treatment except for the device setting (set the device on the volume (V) of the room).

IMPORTANT :

- During diffusion time, leave the room closed and do not enter. The treatment must be conducted with no human presence inside the room.

OXYPHARM

829 rue Marcel Paul
94500 Champigny-sur-Marne
commercial@oxypharm.net
T: +33 1 45 18 78 70

www.oxypharm.net

